

بررسی و مقایسه مورفولوژیکی و فیتوشیمیایی اسانس گیاه دارویی *Echium italicum* L. در رویشگاه‌های مختلف استان کردستان

فرنوش صبوری^{۱*}، براتعلی فاخری^۲، عبدالرحمن رحیمیان بوگر

^۱ کارشناس ارشد گیاهان دارویی، گروه باغبانی، دانشکده کشاورزی، دانشگاه زابل، زابل، ایران

^۲ دانشیار گروه بیوتکنولوژی و اصلاح نباتات، دانشکده کشاورزی، دانشگاه زابل، زابل، ایران

^۳ دانشجوی دکتری گیاهان زیتنی، دانشکده کشاورزی، دانشگاه شیراز، شیراز، ایران

تاریخ دریافت: ۹۵/۱۱/۲۱؛ تاریخ پذیرش: ۹۶/۱۲/۷

چکیده

به منظور بررسی مقایسه ای ویژگی‌های مورفولوژیکی و فیتوشیمیایی اسانس گیاه دارویی گل‌گاوزبان ایتالیایی (*Echium italicum* L.) آزمایشی به صورت طرح آشیانه‌ای در سه رویشگاه سنندج، مریوان و دیواندره استان کردستان در سه ارتفاع مختلف از هر رویشگاه در سال ۱۳۹۳ انجام گرفت. برای اندازه‌گیری شاخص‌های مورد نظر سرشاخه‌های گلدار گیاه از رویشگاه‌های طبیعی در اوایل تابستان برداشت شد. متغیرهای مورفولوژیکی شامل: ارتفاع بوته، وزن تر و خشک، طول و عرض برگ‌های ساده و رزت، فاصله میانگره، قطر ساقه، تعداد گل، قطر ساقه، تعداد برگ و تعداد شاخه فرعی و متغیرهای فیزیولوژیک شامل: شاخص کلروفیل، اندازه‌گیری عناصر NPK خاک، گل و برگ، عملکرد اسانس که با استفاده از روش تقطیر با آب طرح کلونجرا استخراج، شناسایی ترکیبات تشکیل‌دهنده اسانس با استفاده از دستگاه کروماتوگرافی گازی (GC/MS) و عملکرد موسیلاژ نیز با روش جداسازی توسط آب مقطر و رسوب‌دهی در اتانول ۹۶ درصد مورد سنجش قرار گرفتند. داده‌ها به وسیله نرم‌افزار آماری SAS و با روش آنالیز LGM مقایسه میانگین‌ها انجام گرفت. نتایج مقایسه میانگین‌ها نشان داد که با افزایش ارتفاع از سطح دریا و به دنبال آن کاهش درجه، ارتفاع بوته، فاصله میانگره، قطر ساقه، تعداد برگ و درصد موسیلاژ و همچنین اسانس گیاه کاهش، در حالی که شاخص کلروفیل افزایش یافته است. ارتفاع از سطح دریا تاثیر معنی‌داری بر عملکرد اسانس دارد و بالاترین میزان اسانس در پایین‌ترین ارتفاع برای هر رویشگاه بدست آمد، همچنین اکوتیپ سنندج در ارتفاع ۱۵۵۷ متری از نظر عملکرد اسانس (۰/۰۹۷ درصد)، از سایر اکوتیپ‌ها برتر بود. از ترکیبات غالب اسانس می‌توان به بورنئول، کامفور، کانوئیک‌متیل‌استرو کاندینون و... اشاره کرد.

واژه‌های کلیدی: ارتفاع، اکوتیپ، اسانس، رویشگاه، کردستان، گل‌گاوزبان ایتالیایی.

References

- 1-Alipour, N., Mahdavi, K., Mahmoudi, J. and Ghelij-Nia, H. 2015. Investigation into the Effect of Environmental Conditions on the Quality and Quantity of Essential Oil of *Stachys laxa*. Journal of Plant Researches, 28(3): 561-572.
- 2.Assimopoulou, A.N. 2005. Radical scavenging activity of *Alkanna tinctoria* root extract and their main constituent, hydroxyl Naphthoquinones. Phytochemistry Research, 19(2): 7-141.
- 3.Azizi, K., Amini Dehaghi, M. and Abbaspoor, H. 2004. Effect of different air and root-zone temperatures on nitrogen fixation and nodulation of annual medicines. Journal of Agronomy, 3(2): 131-136.
- 4.Chen, H., Zha, J. and Gowans, R.E. 2004. Alcohol enhances HIV type 1 infection in normal human oral keratinocytes by up-regulation cell-surface CXCR4 coreceptor. AIDS Research Hum Retroviruses, 20:513-519.
- 5.Chorli, S., Khorasaninezhad, S., Hemati, Kh. and Kafeshi, B. 2016. Morphological characterization, antioxidant and essential oil content of teas Mountain *Stachys lavandulifolia* Vahl. The habitats of Semnan, Khorasan, 11(41): 41-52.
- 6.Cseke, L.J., Kirakosyan, A., Kaufman, P.B., Warbe, S.L., Duke, J.A. and Briemann, H.L. 2006. Natural Products from Plants 2th Ed. CRC Press, Florida, 569 p.
- 7.Davazdhoami, S. and Majnoon hoseini, N. 2009. Agriculture and manufacturing are some herbs and spices. Tehran University Press, 54p.
- 8.Davise, F.S. and Albrigo L.G. 1994. Citrus. CAB international press, wallington, UK. P, 9814.
- 9.Emad, M., Gheibi, F., Rasuli, M., Khanjanzadeh. and Mohammadi Jozani, S. 2013. Herbalist, borage flower industry. Publications Pune Tehran, 1-48p.
- 10.Firoozjai, M., Hematti, Kh., Khorasannezhad, S., Garmekhkhani A. and Bagherifard, M.A. 2014. The effect of altitude on morphological and phytochemical qualities of nettle leaves (*Urtica dioica* L.) in the provinces of Mazandaran and Golestan. Journal of Plant Ecophysiology Iran, 9(35): 1-11.
- 11.Habibi, H., Mozafarian. N., Majnoon hisseini, N., Chaini, M.R. and Tabatabaee, M.F. 2006. Effect of altitude of essential oil and component in wild Thymus (*Thymus kotschyanus* Bioss) Taleghan vergion. Pazhuhesh and Sazandegi Shahed University, College of Agriculture cept, 73 (In Persian).
- 12.Hasani, J. and Nik Baher, Z. 2014. Evaluation of ecological needs different species of thyme in Kurdistan Habitats. Echo phytochemical Journal of Medicinalplant, 3(1):22-34.
- 13.Hosseinzadeh, H., Shahandeh, S.H. and Shahvand. 2012. Anxiolytic and hypotic effects of aqueous and ethanolic extracts of aerial part of (*Echium italicum* L.) in Mice. Jundishapur Journal Nat Pharm Prod, 7(2): 71-79.
- 14.Jones, K.L. and Muriele, R.R. 1980. Physiological properties of Nitrogen-scavenging Bacteria from the marine environment. Journal of applied Microbiology, 49(3): 421-433.
- 15.Kaviani, B., Khavarinezhad, R. and Fahimi, H. 2005. Interaction phosphorus and potassium on the Kvnlgaysynyn soy protein glycine and β - (*Glycin max* L. Merr) in leaves and callus. Research and development in agriculture and horticulture, 64: 46-57.
- 16.Lotfi, A., Vahabi sadehi, A., Ghanbari, A. and Heidari, M. 2009. The effect of dehydration and manure on the quantity and quality Psyllium (*Plantago ovate Forssk*) in the Sistan region. Journal of Medicinal and Aromatic Plants, (4): 506-518.
- 17.Majd Jabari, T., Rustaian, A. and Vatanpoor, H. 2004. Check ingredient in the essential oil *Tanacetum khorassanicum* (Krasch.) Parsa. Journal of Medicinal and Aromatic Plants, 2(6): 15-20.
- 18.Mamnooi, A. and Sharifi, R. 2011. Effects of water deficit on chlorophyll fluorescence indices and the amount of proline in six barley genotypes and its relation with canopy temperature (canopy) and performance. Plant Biology, 2(5):51-62.
- 19.Malekooni, M., Ramezanpoor, M. and Dastfal, M. 2009. Potassium loss due to drought stress in wheat Darab in Fars. Journal of Soil and Water Sciences, 22(1): 127-135.

20. Meshkani, M., Armin, M. and Jami moini, M. 2013. The response of the physiological and morphological echium (*Borago officinalis*) under biological and chemical fertilizers. Journal of Medicinal Plants, 57(14): 3-61.
21. Mohammadi Soleimani, S. 2005. Influence of environmental factors on essential oil composition of *Teucrium embellished*, rangeland management master's thesis. Light Tarbiat Modarres University.
22. Mohammadi, M., Matinkhah, Kh. and Khajehdin, J. 2014. Some characteristics of ecological Qych (*Zygophyllum artipilicoides*) in some semi-arid regions of Isfahan. Journal of dry canvas, 1(3):1-11.
23. Mohammadian, A., Karamian, R., Mirza, M. and Sepahvand, A. 2015. The effect of altitude and the soil physicochemical properties of the active ingredients of natural habitats Thyme in Lorestan. Scientific Journal Medicinal Plants Research and Perfume, 30(4):519-528.
24. Naderi, M. and Rezai, M. 2005. Phytochemical studies Iranian borage (*Borago officinalis* L.). Iranian Journal of Medicinal and Aromatic Plants Research, 2(3):377-383.
25. Najafpoor, M. 2001. Identify and evaluate the ecological essence of Tehran province. Medicinal and Aromatic Plants Research of Iran. Journal of plant physiology environment, 11(41): 41-52.
26. Noy-Meir, I. 1973. Multivariate analysis of the semi arid vegetation of southern Australia II, Vegetation catenae and environmental gradients. Australian Botanical Journal, 22(1): 40-115.
27. Omidbeigi, R. 2001. The approaches to the production and processing of medicinal plants. Publications Publication Designers, Tehran, (2):46.
28. Page, A.L., Miller, R.H. and Keeney, D.R. 1982. Methods of soil analysis. Part2. chemical precautions and processing. Anim. Waste Manage, 6: 100 – 105.
29. Papageorgios, V.P., Assimopoulou, A.N. and Ballis, A.C. 2008. Alkannins and shikonins: a new class of wound healing agent. Current medicinal Chemistry, 15 (30): 3248-3267.
30. Papagiannis J, Chatzis A. and Sarris G. 2008. Giant right atrial aneurysm: a case report. International Journal Cardiology 129(1): 5-6.
31. Rojcan, I., Dwyer, L.D. and Tollenaar, M. 1999. Note on relationship between leaf soluble carbohydrate and chlorophyll content in maize. Field crop Research, 63:13-17.
32. Saedi, K., Sefidkon, F. and Babai, A. 2015. Phytochemical characteristics of fruit morphology studies and dog rose (*Rosa canina* L.) in northern Iran. Crops Improvement, 16(3):545- 554.
33. Samsam Shariay, H. 2010. Breeding of medicinal plants. Manny Press, 419p.
34. Singh, R.M. 2002. An assessment of ayurvedic concept of cancer and a new paradigm of anticancer treatment in Ayurveda. Journal of Altern complement Maedicinal, 8(5):14-609.
35. Spearing, A.M. and Karlander, E.P. 1979. Effect of high and low temperature on chlorophyll content and metabolism of *Chlorella sorokiniana* shihira and krauss. Environmental and experimental Botany, 19 (4): 237-243.
36. Stainforth, D.A., Aina, T., Christensen, C., Collins, M., Faull, N., Frame, D.J., Kettleborough, J.A., Knight Martin, S.A., Murphy, J.M., Piani, C., Sexton, D., Smith, L.A., Spicer, R.A., Thrope, A.J. and Allen, M.R. 2005. Uncertainty in prediction of the climate response to rising levels of greenhouse gases. Nature, 433(7024):403-406.
37. Xiufeng, Y., Shuangxiu, W., Yang, W., Xinhai, Sh., Shaojun, D. 2004. Soil nutrient factors related to salidroside production of *Rhodiola Chinensis* distributed in Chang Bai Mountain. Journal Environmental and Experimental Botany, 52(2004):267-276.
38. Zargari, A. 2010. Medicinal Plants. Publishing and Printing Institute of Tehran University, 4 p.
39. Zhou, X.M., Madramootoo, G.A., Mackenzie, A.F. and Smith, D.L. 1997. Biomass production and nitrogen uptake in corn-reyegrass systems. Argon Journal, 89(5):749-756.

Effects of ecological factors on morphological and phytochemical characteristics of (*Echium ittalicum* L.) essential oil in Kurdistan natural habitats

Sabouri, F.^{1*}, Fakheri, B.A.², Rahimian-Boogar, A.³

¹Master of Medicinal Plants, Department of Horticulture, College of Agriculture, University of Zabol

²Associate Professor, Department of Biotechnology and Plant Breeding, Faculty of Agriculture, University of Zabol, Zabol, Iran

³PhD student ornamental plants, College of Agriculture, Shiraz University, Iran

Received Time: 9-2-2017 Accepted Time: 26-2-2018

Abstract

In order to investigate the variation of morphological characteristics and essential oil composition of *Echium ittalicum* L., this study was carried out in nested plot design at three different heights with three habitats of Sanandaj, Marivan and Divandareh in Kurdistan province. Aerial parts of plant in flowering stage were collected in summer 2013 from different regions. The plant height, fresh and dry weight, simple and rosette leaf length and width, internodes length, stem diameter, number of flower, number of leaves and branches were evaluated as physiological characteristic and chlorophyll content, measuring elements NPK in the soil, leaves and flowers. Essential oil of plant samples were obtained by Distillation with distilled water (clevenger apparatus) and were analyzed by GS/MS, the mucilage content were evaluated with distill water separation method and sedimentation by 96% ethanol. Results were showed that the shoot height, the internode length, stem diameter, leaf number and percentage of mucilage and essential oil decreased with the altitude increases and so decrease of temperature, while chlorophyll content increases. The borneol, comphene and hexadecanoic acid methyl ester have been higher than other essential oil compounds and the highest essential oil yield obtained in Sanandaj ecotype at 1557 meters above sea level.

Keywords: Ecotype, *Echium ittalicum* L., Habitat, Height, essential oil composition

*Corresponding author; medicinalplantsabouri@gmail.com